

著者	タイトル	雑誌
Hasegawa T, Ushirozako H, Shigeto E, Ohba T, Oba H, Mukaiyama K, Shimizu S, Yamato Y, Ide K, Shibata Y, Ojima T, Takahashi J, Haro H, Matsuyama Y.	<a href="#">The Titanium-Coated PEEK Cage Maintains Better Bone Fusion with the Endplate than the PEEK Cage 6 Months After PLIF Surgery-A Multicenter, Prospective, Randomized Study.</a>	Spine (Phila Pa 1976). 2020 Mar 6. doi: 10.1097/BRS.0000000000003464.
Oba H, Takahashi J, Yokomichi H, Hasegawa T, Ebata S, Mukaiyama K, Ohba T, Ushirozako H, Kuraishi S, Ikegami S, Uehara M, Takizawa T, Munakata R, Hatakenaka T, Matsuyama Y, Haro H.	<a href="#">Weekly Teriparatide versus Bisphosphonate for Bone Union during Six Months after Multi-Level Lumbar Interbody Fusion for Osteoporotic Patients: A Multicenter, Prospective, Randomized Study.</a>	Spine (Phila Pa 1976). 2020 Feb 11. doi: 10.1097/BRS.0000000000003426.
Oba H, Takahashi J, Kobayashi S, Ohba T, Ikegami S, Kuraishi S, Uehara M, Takizawa T, Munakata R, Hatakenaka T, Koseki M, Ebata S, Haro H, Matsuyama Y, Kato H.	<a href="#">Upper instrumented vertebra to the right of the lowest instrumented vertebra as a predictor of an increase in the main thoracic curve after selective posterior fusion for the thoracolumbar/lumbar curve in Lenke type 5C adolescent idiopathic scoliosis: multicenter study on the relationship between fusion area and surgical outcome.</a>	J Neurosurg Spine. 2019 Aug 23:1-8. doi: 10.3171/2019.5.SPINE181469.
Ushirozako H, Hasegawa T, Ebata S, Oba H, Ohba T, Mukaiyama K, Isogai Y, Okada E, Ojima T, Takahashi J, Haro H, Matsuyama Y.	<a href="#">Weekly Teriparatide Administration and Preoperative Anterior Slippage of the Cranial Vertebra Next to Fusion Segment &lt; 2 mm Promote Osseous Union After Posterior Lumbar Interbody Fusion.</a>	Spine (Phila Pa 1976). 2019 Mar 1;44(5):E288-E297. doi: 10.1097/BRS.0000000000002833.
Ebata S, Takahashi J, Hasegawa T, Mukaiyama K, Isogai Y, Ohba T, Shibata Y, Ojima T, Yamagata Z, Matsuyama Y, Haro H.	<a href="#">Role of Weekly Teriparatide Administration in Osseous Union Enhancement within Six Months After Posterior or Transforaminal Lumbar Interbody Fusion for Osteoporosis-Associated Lumbar Degenerative Disorders: A Multicenter, Prospective Randomized Study.</a>	J Bone Joint Surg Am. 2017 Mar 1;99(5):365-372. doi: 10.2106/JBJS.16.00230.